



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

INDEX TO VOLUME 43

New names and final members of new combinations are in **bold face type**

- Abies*, 198; *balsamea*, 354; *lasiocarpa*, 355
Acaena magellanica venulosa, 315; *multifida*, 315; *Philippi*, 315; *platyacantha*, 315
Acantholimon lycopodioides, 581, 583, 640
Acer pennsylvanicum, 244; *rubrum*, 262; *saccharinum*, 260; *spicatum*, 244, 253
Achillea lanata, 272; *lanulosa*, 272, 349; *Millefolium*, 271; *occidentale*, 271; *pannonica*, 272; *subalpina*, 357
Acomastylis sericea, 358; *turbinata*, 357
Aconitum Bakeri, 360; *columbianum*, 349; *delphinifolium*, 353; *heterophyllum*, 582, 632; *insigne*, 360; *Napellus multifidum*, 582, 632; *ramosum*, 363
Acraea rubra, 242, 244
Actinastrum gracillimum, 480
Actinocarya tibetica, 641
Additional species of *Opuntia*, 523
Adenostegia ramosa, 353
Adesmia boronoides, 315; ***campestris***, 315; ***glandulifera***, 315; *parvifolia*, 315; *pumila*, 315; *salicornioides*, 315; *villosa*, 315
Adiantum, 366; *bellum*, 371, 382; *Capillus-Veneris*, 371, 382; *cardiophloea*, 371, 382; *cuneatum*, 365, 366, 382; *pedatum*, 95, 252; *trapeziforme*, 372, 382
Adonis chrysocanthus, 633
Aglaonema, 548
Agropyron andinum, 356; *dentatum elatum*, 628; *Jacquemontii*, 628; *latiglume*, 358; *longe-aristatum*, 628; *repens*, 588, 628; *Scribneri*, 356; *violaceum*, 345
Agrostis alba, 580, 582, 588, 627; *asperifolia*, 349; *canina*, 345, 588, 627; *hiemalis*, 345; *idahoensis*, 358; *variabilis*, 349
Aira caryophyllea, 588, 627
Alisma Plantago, 401
Allardia, 589; *glabra*, 583, 645; *nivea*, 583, 645; *Stoliczkai*, 583, 645; *tomentosa*, 583, 645; *vestita*, 645
Allium, 399, 589; *brevistylum*, 356; *Cepa*, 629; *Govanianum*, 629; *Jacquemonti*, 629; *loratum*, 629; *odorum*, 629; *oreoprasum*, 629; *Pikeanum*, 363; *platyspathum*, 629; *Schoenoprasum*, 629; *Semenovi*, 629; *simillissimum*, 363; *triccoccum*, 252
Allotria, 103; *scabra*, 103
Alnus incana, 247; *rugosa*, 243, 247; *sinuata*, 351
Alopecurus aristulatus, 345; *himalaicus*, 582, 627; *occidentalis*, 345; *pratensis*, 588, 627
Alsine americana, 358; *bailcalensis*, 349; *borealis*, 346; *calycantha*, 349; *crassifolia*, 346; *laeta*, 346; *longifolia*, 346; *polygonoides*, 363
Alsinopsis macrantha, 360; *Nuttallii*, 349
Alstroemeria nana, 310
Alyssum canescens, 586, 634
Amarella anisopetala, 351; *monantha*, 361; *plebeia*, 349; *propinqua*, 354; *strictiflora*, 349
Amaryllis longiflora, 403
Amblystegiella confervoides, 241
Amblystegium fluviatile, 239; *irriguum*, 239; *noterophilum*, 241
Ambrosia trifida, 257, 260
Amphicarpa monoica, 260
Amsinckia angustifolia, 318
Anabaena limnetica, 481; *macrospora*, 482
Anacolia subsessilis, 329
Ananthopus, 103; *clandestinus*, 102; *cordatus*, 105; *undulatus*, 103
Anaphalis contorta, 645; *margaritacea*, 267; *nubigena*, 583, 586, 644; *Stoliczkai*, 645; *subalpina*, 355; *virgata*, 583, 645
Anarthrophyllum Pritchardi, 315
Andreaea petrophila, 241; *Rothii*, 241
Andromeda novae-caesareae, 285, 301; *polifolia*, 347; *tenuinervis*, 302; *Wardiana*, 285, 302
Andropogon Ischaemum, 582, 624
Androsace, 589; *carinata*, 357; *Chamaejasme*, 639; *Chamaejasme coronata*, 639; *filiformis*, 357; *microphylla*, 639; *pinetorum*, 361; *rotundifolia*, 639; *rotundifolia glandulosa*, 639; *sarmentosa*, 639; *sarmentosa primuloides*, 639; *sempervivoides*, 639; *septentrionalis*, 639; *subumbellata*, 357; *villosa*, 581, 583, 624
Aneilema nudiflorum, 96
Anemone albana, 632; *Drummondii*, 353; *globosa*, 349; *lithophila*, 358; *parviflora*, 346; *Richardsonii*, 353; *quinquefolia*, 253; *rivularis*, 632; *rupicola*, 582, 632; *tetonensis*, 358, 359

- Anemonella thalictroides*, 253
Angelica Grayi, 361; *Roseana*, 358, 359
Ankistrodesmus nitzschoides, 473; *setigerus*, 473
Anomobryum filiforme, 68, 328; *procerimum*, 67, 68
Anoetangium Mougeotii, 240
Anomodon attenuatus, 240; *rostratus*, 240
Antennaria acuta, 362; *anaphaloides*, 355; *aprica*, 357; *chilensis*, 320; *chlorantha*, 362; *corymbosa*, 351; *fallax*, 266; *magellanica*, 320; *microphylla*, 357; *media*, 349; *monocephala*, 354; *nana*, 644; *neglecta*, 267; *neodioica*, 266; *petaloidea*, 267; *plantaginifolia*, 265; *plantaginifolia petiolaris*, 265; *rosea*, 349; *sedoides*, 357; *umbrinella*, 355
Anthemis Cotula, 272
Anthiphylla oppositifolia, 347
Anthoceros laevis, 239
Anthracothecium tetraspermum, 148
Anthropogon barbellatum, 361; *elegans*, 357
Anticlea coloradensis, 360
Apocynaceae, 640
Apocynum venetum, 640
Aquilegia columbiana, 351; *elegantula*, 360; *formosa*, 353; *Jonesii*, 358; *larimensis*, 363; *oreophila*, 360; *saximontana*, 360; *scopulorum*, 363; *vulgaris*, 588; *vulgaris pyrenaica*, 582, 632; *vulgaris viscosa*, 632
Arabis alpina, 634; *ambigua*, 353; *Drummondii*, 355; *exilis*, 358; *glabrata*, 634; *glandulosa*, 634; *Lyallii*, 351; *oblanceolata*, 363; *oreophila*, 358, 359; *rupestris*, 355; *tibetica*, 586, 634
Archeria Boltoni, 477
Arachnites uniflora, 311
Aragallus alpicola, 358; *deflexus*, 357; *foliolosus*, 358; *Hallii*, 363; *oreophilus*, 361; *Parryi*, 361; *viscidus*, 358; *viscidulus*, 358
Aralia racemosa, 242
Arctium Lappa, 275, 583, 589, 646; *minus*, 274; *tomentosum*, 275
Arctous erythrocarpa, 353
Arenaria andicola, 314; *foliosa*, 582, 632; *Fendleri*, 360; *Griffithsii*, 582, 632; *holosteoides*, 585, 632; *kashmirica*, 632; *lithophila*, 358; *muscifformis*, 585, 632; *norvegica*, 349; *salmonensis*, 353; *serpyllifolia*, 588, 632; *serpylloides*, 314; *Stracheyi*, 632; *Tweedyi*, 360
Argentina subarctica, 354
Arjona pusilla, 313; *tuberosa*, 313
Arisaema quinatum, 95; *triphyllum*, 242
Armeria chilensis, 317
Arnebia tibetana, 641; *Thomsoni*, 641
Arnica cordifolia, 349; *macilenta*, 362; *ovata*, 362; *Parryi*, 355; *pedunculata*, 349; *pumila*, 357; *rhizomata*, 357; *Rydbergii*, 355; *subplumosa*, 357; *tenuis*, 363
Artemisia Absinthium, 274, 646; *amygdalina*, 583, 646; *annua*, 273; *biennis*, 273, 645; *caudata*, 273; *desertorum*, 645; *Dracunculus*, 645; *Falconeri*, 645; *laciniata*, 645; *macrocephala*, 646; *maritima*, 645; *minor*, 586, 646; *parviflora*, 645; *Pattersonii*, 362; *persica*, 645; *pontica*, 274; *Richardsoniana*, 354; *sacrorum*, 645; *salsoloides*, 583, 645; *saxicola*, 363; *scopulorum*, 357; *scoparia*, 645; *Siversiana*, 646; *spithamea*, 346; *stelleriana*, 274; *Stracheyi*, 646; *stricta*, 645; *Tournefortiana*, 645; *vulgaris*, 273
Arthonia atrata, 149; *conferta*, 149; *polymorpha*, 149; *rubella*, 149
Arthothelium spectabile, 149
Asarum canadense, 242, 253
Asclepiadaceae, 640
Asperugo procumbens, 641
Aspidium mohrioides, 309; *noveboracense*, 252; *spinulosum*, 242
Asplenium acrostichoides, 242; *angustifolium*, 95; *bulbiferum*, 376, 382; *Filix-femina*, 261, 373, 382; *magellanicum*, 309; *platyneuron*, 375, 382; *pycnocarpon*, 95; *Trichomanes*, 242
Aster acuminatus, 244; *alpinus*, 357; *apricus*, 355; *Canbyi*, 362; *Fremontii*, 357; *heterochaeta*, 583, 586, 644; *paniculatus*, 261; *tibeticus*, 581, 586, 644; *Vahlilii*, 321
Astragalus adesmiaefolius, 636; *alpinus*, 587; *brevicaulus*, 315; *cicerifolius*, 637; *ciliolatus*, 636; *coluteocarpus*, 582, 637; *confertus*, 636; *Dusenii*, 315; *Falconeri*, 636; *frigidus*, 581, 582, 637; *Heydei*, 636; *himalayanus*, 582, 636; *hypoglottoides*, 637; *macropterus*, 580, 636; *multiceps*, 637; *Munroi*, 580, 637; *melanostachys*, 582, 636; *nivalis*, 580, 637; *ophiocarpus*, 636; *oxyodon*, 636; *patagonicus*, 316; *peduncularis*, 637; *rhizanthus*, 582, 637; *Royleanus*, 637; *strictus*, 636; *subulatus*, 583, 637; *tibetanus*, 582, 636; *tribulifolius*, 636; *Webbianus*, 637; *zanskarensis*, 637
Atelanthera perpusilla, 634
Atelephragma elegans, 357; *debilis*, 358; *Forwoodii*, 358, 359
Atriplex crassifolia, 580, 585, 631; *hortensis*, 631; *rosea*, 585, 588, 631
Avena fatua, 548, 554, 588, 627; *Mortoniana*, 360; *sativa*, 637; *striata*, 345
Axyris amaranthoides, 582, 631; *amaranthoides humifusa*, 631
Azaliastrum albiflorum, 351
Azorella trifoliata, 317; *trifurcata*, 317
Baccharis Darwinii, 319; *halmifolia*, 265;

- magellanica, 319; patagonica, 319; **Shaferi**, 461
Bacidia fusciorubella, 153
 Balsaminaceae, 638
Barbarea vulgaris, 587
Barbula costata, 325; *Eubryum*, 66; *exesa*, 67; *pruinosa*, 325; *replicata*, 325; *subulatula*, 326
 BARTHOLOMEW, E. T. Observations on the fern rust *Hyalopsora Polypodii*, 195
Bartramia pomiformis, 240, 308
Bauhinia gigantea, 295; **Jenningsii**, 463; *marylandicus*, 295; **ripleyensis**, 285, 294
Bazzania trilobata, 244
 BENEDICT, R. C. The origin of new varieties of *Nephrolepis* by orthogenetic saltation. I. Progressive variations, 207
Benzoin aestivale, 243
 Berberidaceae, 314, 633
Berberis empetrifolia, 314; *ilicifolia*, 314; *microphylla*, 314; *ulicina*, 633; *vulgaris*, 588, 632
 Bermuda, The lichens of, 145
 BERRY, E. W. Contributions to the Mesozoic flora of the Atlantic coastal plain,—XI. Tennessee, 283
Besseyia alpina, 361; *gymnocarpa*, 357; *plantaginea*, 361; *reflexa*, 363; *Ritteriana*, 362
Betula alba papyrifera, 252; *glandulosa*, 345; *utilis*, 578, 630
 Betulaceae, 630
Biatora fusciorubescens, 152
 BICKNELL, E. P. The ferns and flowering plants of Nantucket—XVII, 265
Bidens cernua, 269; *cernua elliptica*, 269; *connata*, 269; *frondosa*, 270; *petiolata*, 269; *vulgata*, 257, 270
Biebersteinia Emodii, 586, 637
Bilimbia Brittoniana, 152; **floridana**, 152; **sphaeroides vacillans**, 153
Bistorta bistortoides, 349; *linearifolia*, 356; *vivipara*, 346
Blastenia floridana, 159
Blephariglottis conspicua, 408
Blumea bifoliata, 644
Boehmeria cylindrica, 261
 Boraginaceae, 318, 641
 Botanical work of Dr. E. W. Hilgard, 389
Botrychium Lunaria, 309; *virginianum*, 252, 257
Brachyactis robusta, 644; *umbrosa*, 644
Brachythecium asperulum, 333; *flaccum*, 333; *rivulare*, 239; *plumosum*, 241; *sericeovirens*, 333; *stereopoma*, 334
 Branched prothallia in the Polypodiaceae, 365
Brassica juncea, 588; *Napus*, 634; *nigra*, 588
Braunia canescens, 330; *Elliottii*, 70
Braya alpina, 586, 634; *humilis*, 587; *rosea*, 634; *tibetica*, 634; *uniflora*, 634
Breutelia bryocarpa, 329; *tomentosa*, 330
 BRITTON, N. L. Studies of West Indian plants—VIII, 441
Briza media, 588, 627
Bromus altissimus, 261; *barbatus*, 585, 628; *ciliatus*, 345; *coloratus*, 309; *crinitus*, 585, 627; *macrostachys*, 585, 628; *oxyodon*, 628; *patulus*, 627; *polyanthus*, 349; *tectorum*, 580, 588, 627
 Bryaceae, 625
Bryum andicola, 328; *argenteum*, 328; **biforme**, 328; *canariense*, 69; **canariensisiforme**, 68; *densifolium*, 329; *polytrichoides*, 69; *procerrimum*, 68; *promontorii*, 68; *provinciale*, 69; *tibeticum*, 582, 625
Buellia canescens, 158; *myriocarpa*, 159; *parasema*, 159
Bulbostylis alpestris, 446; *capillaris*, 447; *fimbriata*, 446; *floccosa*, 445; *leucostachys*, 445; *paradoxa*, 444; *subaphylla*, 446; *Tuerckheimii*, 446
Bupleurum falcatum, 583, 639; *falcatum nigrocarpa*, 581, 639; *tenuis*, 639
Byrsonima lucida, 456; **ophiticola**, 456
 Caesalpiniaceae, 294
Caesoma interstitiale, 559; *nitans*, 559–569
Calamagrostis canadensis, 345; *Langsdorfi*, 345; *littorea*, 627; *purpurascens*, 345; *Stoliczkae*, 627
Calamintha umbrosa, 642
Calceolaria biflora, 318; **Pritchardi**, 318; *tenella*, 318; *uniflora*, 318
Callymenia reniformis, 171, 173
Caltha leptosepala, 358; *rotundifolia*, 360
 Calyceraceae, 319
Calyptanthus Boldinghi, 466; *Maxoni*, 466; *nummularia*, 466; **pinetorum**, 465
 Cambial activity in certain horticultural plants, 533
Campanula, 589; *argyotricha*, 644; *aristata*, 583, 644; *colorata tibetica*, 644; *cashmiriana*, 644; *heterodoxa*, 354; *latifolia*, 583, 644; *Parryi*, 362; *petiolata*, 346
 Campanulaceae, 644
Camptosorus, 366; *rhizophyllus*, 372, 382
Campylopus aethiops, 65; **angolensis**, 65; *flexuosus*, 65; *introflexus*, 65; *leucogonodes*, 324
Capnorchis brachycarpum, 363
 Capparidaceae, 634
Capparis spinosa, 634
 Caprifoliaceae, 644
Capsella elliptica, 634; *Thomsoni*, 634
Caragana cuneata, 636; *pygmaea*, 581, 586, 636
Cardamine acuminata, 355; *hirsuta*, 314;

- macrophylla, 634; pratensis, 587; umbellata, 349
Carduus nutans, 589, 646; *Thomsoni*, 646
Carex adusta, 616, 617; **abrupta**, 604, 618, 620; *aenea*, 347; *albonigra*, 356; **amplectens**, 603, 611; *athrostachys*, 349, 603; *atrata*, 345; *atrostiformis*, 347; *aurea*, 345; *Banksii*, 310; *bella*, 360; *brunnescens*, 345; *canescens*, 345; *capillaris*, 346; *capitata*, 347; *chalciolepis*, 356; *chimaphila*, 356; **convoluta**, 424, 428; **Davyi**, 602, 606, 615; *deflexa*, 347; *diluta*, 628; *Douglasii*, 349; *eburnea*, 360; *Egglestonii*, 360; *ely-noides*, 360; *Engelmanii*, 355; *festiva*, 611; *festivella*, 349; *feta*, 602; *Geyeri*, 355; *glaucescens*, 407; *Goodenovii*, 587; **gracilior**, 603, 614; *gynocrates*, 345; *haematostoma*, 628; *Halleri*, 346; *Hassei*, 347; **Harfordii**, 604, 615; *Helio-mastes*, 347; *Hepburnii*, 345; *hirtella*, 628; *illota*, 349; *inconspicua*, 310; *incurva*, 628; **integra**, 604, 608; *interior*, 345; *laeviculmis*, 351; **lancifructus**, 602, 607; *leporina*, 602; **leporinella**, 602, 605; *leptalea*, 345; *livida*, 347; *MacCallae*, 353; **mariposana**, 604, 619; *microglochin*, 628; *microptera*, 603; *Mertensii*, 351; *miliaris*, 347; *misandra*, 346; *Moorcroftii*, 585, 628; **multicostata**, 602, 604; *multimoda*, 611; *neglecta*, 425; *Nelsonii*, 360; *nigricans*, 349; *nivalis*, 584, 628; *nova*, 349; *obtusata*, 345; *occidentalis*, 360; *oligocarpa*, 628; **olympica**, 603, 610; **pachycarpa**, 604, 616; *pachystachya*, 611; *parva*, 628; **paucifructus**, 604, 615; *petasata*, 355, 606; *phaeocephala*, 349, 602; *physocarpa*, 353; *praticola*, 602; *Preslii*, 611; *pseudoscupoidea*, 356; *pyrenaica*, 355; *radiata*, 425, 432; *Reynoldsii*, 349; *rigida*, 582, 587, 628; *rosea*, 423, 435; *rosea radiata*, 423; *rosea minor*, 423; *Rossii*, 345; *rostrata*, 346; *scirpoidea*, 347; *siccata*, 345; *stellulata*, 424; *stenochlaena*, 351; *stenophylla*, 587, 628; *straminiformis*, 602; **sub-bracteata**, 603, 612; *subfusca*, 603, 608, 609; *supina*, 347; **teneraeformis**, 603, 609; *tenuirostris*, 603, 605; *Tolmiei*, 351; *trisperma*, 424; *vaginata*, 347; *venustula*, 353; *verrucosum*, 95, 407; *vulgaris*, 628
Carex, Notes on, —X, 423; —XI, 601
Carex rosea and its allies, 423
Carpinus caroliniana, 243, 252
Carum Carvi, 588, 639
Carya cordifolia, 262; *cordiformis*, 252
Caryophyllaceae, 3, 13, 631
Casasia parvifolia, 461
Cassia bahamehsis, 459; *domingensis*, 459; *nictitans*, 463; *procumbens*, 463; *pygmaea*, 463; **scleroxyla**, 458; **Shaferi**, 459
Cassine magellanica, 316
Cassiope Mertensiana, 352; *saximontana*, 362; *tetragoma*, 347
Castilleja brunnescens, 362; *confusa*, 362; *lanata*, 357; *lancifolia*, 357; *lutea*, 351; *occidentalis*, 357; *pallidescens*, 351; *pallida*, 354; *puberula*, 362; *pulchella*, 359; *rhexifolia*, 357; *Tweedyi*, 359
Catabrosa aquatica, 345, 587, 627; *Thomsoni*, 627
Catharinaea undulata, 239, 246
Caulophyllum thalictroides, 242, 253
Celastraceae, 316
Centaurea Cyanus, 276; *depressa*, 646; *melitensis*, 276
Cephaloziella Hampeana, 240
Cerastium arvense, 313; *beeringianum*, 349; *Earlei*, 360; *oreophilum*, 360; *scopulorum*, 356; *strictum*, 356; *trigynum*, 632; *variabile*, 360; *vulgatum*, 588, 632; *vulgatum nepalense*, 632; *vulgatum tibetica*, 632
Cestrum pinetorum, 461
Chaerophyllum acuminatum, 583, 639; *villosum*, 639
Chaetosphaeridium globosum, 471; **ovalis**, 471; *Pringsheimii*, 471
Chamaecrista micrantha, 463; **savannarum**, 463
Chamaerhodos subulosa, 636
Chamaesyce Cowellii, 457; **Jenningsii**, 465
Characium stipitatum, 472
Cheiranthus albiflorus, 634; *himalayensis*, 634; *Stewartii*, 634
Cheirinia amoena, 360; *nivalis*, 360; *oblanceolata*, 360; *radicata*, 360
Chenolea divaricata, 631
Chenopodiaceae, 313, 631
Chenopodium album, 313, 585, 588, 631; *antarcticum*, 313; *Botrys*, 588, 631; *Blitum*, 631; *glaucum*, 588, 631; *hybridum*, 587, 631
Chiloscyphus rivularis, 239
Chimaphila umbellata, 246
Chiodecton Montagnei, 151
Chiogenes hispidula, 347
Chionophila Jamesii, 361
Chlamydomonas stipitata, 472
Chloraea Kingii, 312; *magellanica*, 312; *pleistodactyla*, 312
Chloris virgata, 627
Chodatella ciliata, 477; *ciliata minor*, 477; *subsalsa*, 477
Chondriella graminea, 647
Chorisporea sabulosa, 582, 634
Christolea crassifolia, 580, 586, 634
Chroococcus limneticus, 481; *limneticus distans*, 481
Chrysanthemum Balsamita, 273; *Leu-*

- canthemum, 272; *Leucanthemum pin-*
natifidum, 272; *Parthenium*, 273;
Parthenium breviradiatum, 273;
Richteria, 581, 583, 645; *Stoliczkai*,
 645; *tibeticum*, 645
Chrysopsis alpicola, 362; *asprella*, 362;
pumila, 357
Chrysosplenium americanum, 242
 CHURCH, M. B. The development of the
 embryo sac and embryo of *Cooperia*
Drummondii, 397
Cibotium Chamissoi, 387; *Menziesii*, 387
Cicer songaricum, 583, 637
Cinna arundinacea, 261
Cinnamomum Heeri, 285, 298; sp., 285
Circaea alpina, 242; *lutetiana*, 261
Cirsium arvense, 276; *discolor*, 275;
horridulum, 275; *lanceolatum*, 275;
odoratum, 275
Cissites crispus, 285, 296; *Newberryi*, 296
Cladonia coccifera, 308; *fimbriata bar-*
bonica, 153; *fimbriata nemoxyna*, 153;
fimbriata simplex, 153; *mitrula*, 154;
pityrea, 154; *rangiformis pungens*, 154
Claytonia megarrhiza, 356; *virginica*, 253
Clematis orientalis, 590, 632; *rhodantha*,
 356
Climacium americanum, 239
Clitoria glomerata, 452
Cnicus arvensis, 580, 646; *Wallichii*, 646
Cochlearia scapiflora, 633
Codonopsis ovata, 644
Codonorchis Lessonii, 312
Coelosphaerium Naegelianum, 472
Coleosanthus umbellatus, 362
Collema bermudanum, 155; *flaccidum*,
 155; *furvum*, 155; *nigrescens*, 155;
thamnodes, 155
 COLLINS, F. S. & HOWE, M. A. Notes
 on species of *Halymenia*, 169
Collinsonia canadensis, 242
Collomia coccinea, 318; *gracilis*, 318;
debilis, 358, 359
Cololejeunea Biddlecomiae, 241
Colutea arborescens, 636
Comarum palustre, 347
Commelina angustifolia, 97, 105, 108, 407;
caroliniana, 97, 99, 407; *communis*, 97,
 98, 407; *crispa*, 97, 107, 407; *dianthi-*
folia, 97, 102; *diffusa*, 100; *elegans*, 97,
 110, 407; *erecta*, 97, 104, 105, 407;
hirtella, 102, 407; *ignorata*, 103;
longicaulis, 97, 100, 407; *longifolia*,
 102; *Nashii*, 109; *nudiflora*, 96, 407;
ignorata, 103; *Rafinesquei*, 105; *saxi-*
cola, 105; *Swingleana*, 108; *virginica*,
 97, 102, 407
Commelina (Plumier) L. in the United
 States, The genus, 96
 Compositae, 319, 644
Condrophora americana, 361; *Fremontii*,
 361
Condrosea Aizoon, 348
Coniomitella Williamsii, 358
Conium maculatum, 317
 Connecticut, The vegetation of, 235
Conocephalum conicum, 239, 246
Conringia planisiliqua, 634
 Contributions to the Mesozoic flora of
 the Atlantic coastal plain—XI. Ten-
 nessee, 283
 Convolvulaceae, 640
Convolvulus arvensis, 588, 640
Cooperia Drummondii, The development
 of the embryo sac and embryo of, 397
Coprosmanthus herbaceus, 414; *peduncu-*
laris, 414; *tenuifolius*, 411
Cordia Sebestena brachycalyx, 457
Corispermum hyssopifolium, 587, 631
Coronopus australis, 314
Cortaderia pilosa, 309
Corydalis adiantifolia, 633; *crassifolia*,
 586, 633; *flabellata*, 580, 633; *Gortscha-*
kovii, 582, 633; *meifolia*, 584, 633;
Moorcroftiana, 633; *stricta*, 586, 633;
tibetica, 633
Corylus americana, 246
Cotoneaster acuminata, 635; *nummularia*,
 635; *vulgaris*, 635
Cousinia Falconeri, 581, 646
Crambe cordifolia, 634
 Crassulaceae, 635
Cratoneuron filicinum, 241
Cremanthodium Decaisnei, 583, 646;
oblongatum, 646
Crepis flexuosa, 647; *Stoliczkai*, 647;
tenuifolia, 647
Crescentia portoricensis, 457
 Cruciferae, 314, 633
Cryptotaenia canadensis, 242
Cryptocarpus Curtisi, 117
Ctenidium malacodes, 333
Culcitium magellanicum, 320
Cuscuta capitata, 583, 640; *europaea*, 588,
 640
Cyclodictyon aeruginosum, 332
Cylindrospermum stagnale angustatum,
 481
Cynanchum acutum, 581, 640; *glaucum*,
 640; *Heydei*, 640
Cynoglossum petiolatum, 641; *Wallichii*,
 641
 Cyperaceae, 310, 628
Cyperus cyperoides, 441; *fugax*, 441;
granularis, 441; *Haspan*, 462; *nanus*,
 441, 442; *ignotus*, 442; *pinetorum*,
 462
Cypripedium passerinum, 347
Cystium platytropus, 358
Cystopteris bulbifera, 242; *fragilis*, 195,
 242, 309, 582, 587, 625
Cytherea bulbosa, 346; *occidentalis*, 353
Dactylis glomerata, 588, 627

- Danthonia californica*, 349; *kashmiriana*, 627
Dasiphora fruticosa, 346
Dasystephana affinis, 357; *calycosa*, 353; *glauca*, 353, 354; *monticola*, 358; *Parryi*, 361; *Romanzovii*, 357
Delesseria sinuosa, 180
Delphinium attenuatum, 363; *Barbeyi*, 360; *Brunonianum*, 581, 582, 584, 585, 632; *cashmirianum*, 582, 632; *occidentale*, 360
Dentaria diphylla, 253; *laciniata*, 253; *rupicola*, 353
Deschampsia atropurpurea, 345; *caespitosa*, 345, 585, 587, 627; *curtifolia*, 356; *flexuosa*, 309; *koelerioides*, 627
Desmopsis, 184; *bibracteata*, 190, 192, 193; *Galeottiana*, 185-188; *Maxonii*, 185, 188, 193; *panamensis*, 185, 189, 192, 193; *Oerstedii*, 191
Desmopsis, A new genus of *Annonaceae*, 183
Desmos, 183, 185; *chinensis*, 183, 184; *cochinchinensis*, 183
Dewalquea insignis, 294; *Smithi*, 285, 293
Development of the embryo sac and embryo of *Cooperia Drummondii*, The, 397
Development of the embryo and seedling of *Dioscorea villosa*, The, 545
Deyeuxia compacta, 627; *scabrescens*, 627
Dianthus anatolicus, 582, 584, 632; *angulatus*, 632; *crinitus*, 632; *deltoides*, 588, 632; *Seguieri*, 632
Dicentra cucullaria, 253
Dichotomaria, 621
Dicranum fulvum, 241; *Mittenii*, 323
Didymodon afer, 66; *rubellus*, 240
Dilophia salsa, 633
Dioclea galactioides, 451; *rudolphioides*, 449
Diodia arenicola, 467; *ciliata*, 467
Diospyros primaeva, 285; *virginica*, 303
Dioscorea bonariensis, 546; *brasiliensis*, 546; *cordifolia*, 546; *paniculata*, 545; *pyrenaica*, 546, 555; *sinuosa*, 546; *villosa*, 545
Dioscorea villosa, The development of the embryo and seedling of, 545
Discaria discolor, 316
Disporum trachycarpum, 356
Distegia involucrata, 345
Dixon, H. N. New and rare African mosses, from Mitten's herbarium and other sources, 63
Dodecatheon uniflorum, 363
Doronicum Falconeri, 646
Douglasia montana, 358; *nivalis*, 362
Draba alpina, 582, 634; *andina*, 356; *aurea*, 346; *aureiformis*, 361; *brachystylis*, 363; *chrysantha*, 361; *crassa*, 356; *crassifolia*, 346; *densiflora*, 358; *eurocarpa*, 353; *fladnizensis*, 346; *glacialis*, 582, 634; *graminea*, 361; *incana*, 634; *lasiophylla*, 584, 634; *lonchocarpa*, 351; *luteola*, 361; *nitida*, 349; *oligosperma*, 349; *Parryi*, 360; *pectinata*, 350; *spectabilis*, 361; *sobolifera*, 363; *streptocarpa*, 361; *tibetica*, 634; *ventrosa*, 363
Dracocephalum heterophyllum, 586, 642; *nutans*, 583, 642; *stamineum*, 642
Drepanocladus aduncus, 77; **Hallii**, 77; **longifolius**, 332
Drimys Winteri, 314
Drosera longifolia, 2, 3, 348; *rotundifolia*, 1-27, 347; *Whittakeri*, 3
Drosera rotundifolia, Physiological observations on, 1
Drymocallis fissa, 361; *pseudorupestris*, 351; *pumila*, 350
Dryophyllum gracile, 285, 290
Dugaldia Hoopesii, 349
Duthiea bromioides, 628

Ebenaceae, 303
Echinops cornigerus, 646; *niveus*, 646
Echinosperrum barbatum, 641; *Redowskii*, 641; *semiglabrum*, 641
Edwinia macrocalyx, 363
Elaeagnaceae, 638
Elaeagnus hortensis, 638
Eleocharis microcarpa, 442; *nodulosus*, 442; *paulstris*, 587, 628
Elephantella groenlandica, 346
Elephantopus arenarius, 468; *pratensis*, 468
Elodea canadensis, 255
Elsholtzia cristata, 642; *densa*, 583, 642; *pusilla*, 584, 642
Elymus canadensis, 261; *dahuricus*, 582, 628; *dasystachys*, 628; *glaucus*, 349; *sibiricus*, 628; *simplex*, 360
Embothrium coccineum, 313
Encalypta contorta, 241
Endophyllum, 566; *Sempervivi*, 566
Endymion nutans, 400
Entodon brevirameus, 72; *Dregeanus*, 72; *platygyrioides*, 331
Ephedra Gerardiana, 580, 582, 626; *intermedia*, 626
Epilobium alpinum, 346; *anagallidifolium*, 346; *angustifolium*, 583, 587, 638; *clavatum*, 357; *Drummondii*, 355; *Hornemannii*, 346; *latifolium*, 583, 638; *latiusculum*, 358; *minutiflorum*, 638; *ovatifolium*, 361; *patagonicum*, 317; *Royleanum*, 581, 638; *stramineum*, 361; *wyomingense*, 357
Epipactis latifolia, 629
Equisetaceae, 626
Equisetum arvense, 257, 587, 626; *diffusum*, 626; *fluvatile*, 255; *ramosissimum*, 626

- Eragrostis minor*, 588
Erechtites hieracifolia, 274
Eremurus himalaicus, 629
Erianthus Ravennae, 626
 Ericaceae, 301
Erigeron alpinus, 320, 347, 644; *alpinus* *multicaulis*, 583; *alpinus uniflorus*, 583, 644; *andryaloides*, 581; *bellioides*, 644; *compositus*, 346; *Coulteri*, 349; *elator*, 362; *flabelliformis*, 363; *flagellaris*, 357; *formosissimum*, 362; *glabellus*, 357; *glandulosus*, 362; *grandiflorum*, 355; *jucundus*, 346; *Kingii*, 350; *leiomeris*, 362; *lanatus*, 354; *melanocephalus*, 362; *multicaulis*, 644; *multiradiatus*, 644; *pedatus*, 351; *pinnatisectus*, 362; *pulchellus*, 246; *purpureum*, 347; **purpuripes**, 468; *radicatus*, 359; *superbus*, 362; *spiculosus*, 320; *trifidus*, 357; *unalaschkensis*, 347
Eriocaulon decangulare, 95; *lineare*, 390; *septangulare*, 390
Eriogonum arcuatum, 360; *caespitosum*, 358; *chloranthum*, 360; *flavum*, 356; *heracleoides*, 358, 359; *neglectum*, 360; *Piperi*, 351; *polyphyllum*, 351; *pyrolae-folium*, 353; *umbellatum*, 349
Eriophorum alpinum, 348; *angustifolium*, 345; *Chamissonis*, 347; *Scheucheri*, 348
Eritrichium strictum, 581, 641; *spathulatum*, 641; *tibeticum*, 641
Erodium cicutarium, 316, 588, 637; *Stephanianum*, 637; *tibetanum*, 637
Erpodium grossirete, 70; *Hanningtonii*, 69; *Joannis-Meyeri*, 69; *Menyhardtii*, 69
Ervum lens, 637
Eryngium Billardieri, 639
Erysimum hieracifolium, 634; *odoratum*, 634
Erythrodontium squarrosum, 331
Erythronium, 399; *americanum*, 252; *grandiflorum*, 351; *obtusum*, 358; *parviflorum*, 360
Escallonia Britteniana, 314; *rubra*, 314
Euclidium syriacum, 634
Eugenia anceps, 285, 301
Eupatorium articaefolium, 261; *purpureum*, 261
Euphorbia Maddenii, 638; *portulacoides*, 316; *Thomsoniana*, 638; *tibetica*, 638
 Euphorbiaceae, 295, 316, 638
Euphrasia mollis, 354; *officinalis*, 643
Eurotia ceratioides, 580, 585, 631
Evolvulus arenicola, 466

Fabronia, 73; *andina*, 332; *polycarpa*, 332; **Wageri**, 73
 Fagaceae, 297, 313
Fagopyrum esculentum, 631; *tartaricum*, 631
 Ferns and flowering plants of Nantucket, The, —XVII, 265
 Fertilization in *Fritillaria pudica*, 505
Ferula Jaeschkeana, 639
Festuca, 589; *altaica*, 627; *arizonica*, 360; *crassipes*, 285, 291; *curtifolia*, 345; *gracillima*, 309; *minutiflora*, 360; *Myuros*, 588, 627; *nitidula*, 627; *purpurascens*, 310; *rubra*, 585, 587, 627; *sibirica*, 627; *Thurberi*, 360
Ficus atavina, 291; *crassipes*, 285, 291; *Krausiana*, 285, 291; *Mohliana*, 292; *ovatifolia*, 285, 293; *Woolsoni*, 293
Filago arvensis, 644
Fimbristylis miliacea, 443
Fissidens adiantoides, 239; *asplenioides*, 324; *crispus*, 324; *cristatus*, 241; *incurvus*, 239; *rigidulus*, 324; *subobtusatus*, 65
 Flora of Ladak, Western Tibet, The, —I. Discussion of the flora, 571; —II. Check list of Ladak plants, 625
Fontinalis, 239; *dalecarlica*, 239; *gigantea*, 239; *Lescurii*, 239
Fragaria bracteata, 349; *glaucia*, 357
Fraxinus americana, 260, 263; *excelsior*, 640; *nigra*, 263; *pennsylvanica*, 260, 263
Fritillaria pudica, 505; *Roylei*, 629; *tenella*, 505
Frullania Asagrayana, 240; *riparia*, 241
Fuchsia magellanica, 317
Funaria calvescens, 328; *submicrostoma*, 580, 625
 Fumariaceae, 625
 Furlong, Charles Wellington, Plants from southern Patagonia collected by, 305
 Further records of Jamaica sedges, 441
 Further studies of the orange rusts of *Rubus* in the United States, 559

Gagea persica, 629
Galactia angustifolia, 450; *Berteriana*, 450; *brachyodon*, 449, 451; *Combsii*, 448, 450; *cubensis*, 450; **Earlei**, 448, 450; *galactioides*, 449, 451; *Grisebachii*, 449; *impressa*, 451; **Jenningsii**, 449, 451; *Jussiaeana*, 449, 452; *minutifolia*, 448, 449; *monophylla*, 448, 449; *Odonia*, 449; *parvifolia*, 448, 449; *revoluta*, 449, 451; *rudolphioides*, 448, 449; **savannarum**, 449, 452; *spiciformis*, 448, 450; *stenophylla*, 449; *striata*, 448, 450; **suberecta**, 448, 450
Galactia P. Br. in Cuba, 448
Galaxaura obtusata, A note on the structural dimorphism of sexual and tetrasporic plants of, 621
Galinsoga aristulata, 270; *brachystephana*, 270; *caracasana*, 271; *hispida*, 270, *Humboldtii*, 271; *parviflora*, 270, 271; *parviflora hispida*, 270; *quadriradiata*, 271; *quinquiradiata*, 271; *urticaefolia*, 271
Galium Aparine, 319, 587, 643; *boreale*,

- 583, 587, 643; pauciflorum, 586, 643; tricorne, 643; verum, 583, 589, 643
Gaultheria humifusa, 348; *ovatifolia*, 351
Gayophytum racemosa, 349
Gentiana aquatica, 586, 640; *aurea*, 583, 640; *borealis*, 640; *carinata*, 583, 640; *decumbens*, 583, 640; *Hugelii*, 640; *humilis*, 581, 640; *leucomelaena*, 640; *Moorcroftiana*, 583, 640; *patagonica*, 318; *pseudo-humilis*, 640; *pygmaea*, 640; *serrata*, 581, 640; *serrata* **Stracheyi**, 640; *squarrosa*, 640; *tenella*, 583, 640; *Thomsoni*, 640
 Gentianaceae, 318, 640
 Genus *Machaonia* H. & B. in Cuba, The, 452
Georgia pellucida, 246
 Geraniaceae, 637
Geranium collinum, 583, 637; *grandiflorum*, 637; *maculatum*, 246; *magellanicum*, 316; *Pattersonii*, 361; *polyanthes*, 637; *pratense*, 637; *sessiliflorum*, 316; *sibiricum*, 637
Gerardia pinetorum, 466
Geum magellanicum, 315; *rivale*, 346
Gifola germanica, 589, 644
Gilia globularis, 361
Glaux maritima, 587, 639
Globulina peruviana, 325
Gloeocystopsis, 474; **limneticus**, 475
Gloeocystis Rehmanni, 475
Glyceria distans, 627; *poaeoides*, 627; *Thomsoni*, 627
Glyphis cicatricosa, 151
Glyphomitrium incurvum, 241
Gnaphalium obtusifolium, 267; *purpureum*, 268, 320; *spicatum*, 320; *Stewartii*, 645; *uliginosum*, 257, 268
 Gnetaceae, 626
Golenkinia botryoides, 477; *fenestrata*, 477
Gomphosphaeria Naegeliana, 472
Gormannia debilis, 363
 Gramineae, 309, 626
Graphis Afzelii, 150; *Lineola*, 150; *Pavoniana*, 150; *scripta*, 151; *scripta serpentina*, 151; *striatula*, 151
Grapphephorum muticum, 349; *Wolfii*, 360
 Griffiths, D., Additional species of *Opuntia*, 523; New species of *Opuntia*, 83
Grimaldia fragrans, 241
Grimmia apocarpa, 239; *conferta*, 240; *micro-ovata*, 327; *ovata*, 327; **rivulariopsis**, 327
Gyalecta Farlowi, 145, 152; *hyalina*, 152
Gymnoconia interstitialis, 559-569
Gymno-ischyrodon, 73
Gymnostomum rupestre, 244
Gypsophila, 632
Gyrostomum scyphuliferum, 151
Gyrotheca tinctoria, 408
Habranthus gracilifolius Borthianus, 398
Haematomma puniceum, 158
Halarachnion ligulatum, 173
Halenia elliptica, 640
 Haloragidaceae, 638
Halocharis violaceae, 631
Haloxylon Thomsoni, 631
Halymenia actinophysa, 169, 180, 181, 182; **bermudensis**, 169, 173, 180, 182; **echinophysa**, 180, 181; *Floresia*, 169-179; *floridana*, 169-176, 182; *floridana dentata*, 177; **Gelinaria**, 173, 175-177, 179, 182; **pseudofloresia**, 177-179, 182
Halymenia, Notes on species of, 169
Halymenites major, 285, 303
Hamamelis virginiana, 243
Haplocladium amplexicaule, 76; *angustifolium*, 76; *afro-capillatum*, 76; *capillatum*, 76
Haplohymenium triste, 240
 HARPER, R. M. Botanical work of Dr. E. W. Hilgard, 389
 HARRIS, J. A. A quantitative study of the factors influencing the weight of the bean seed—II. Correlation between number of pods per plant and seed weight, 485
Hedwigia albicans, 240, 330; *imberbe*, 330
Hedysarum Mackenzii, 354; *microcalyx*, 583, 637; *sulphurescens*, 351
Helenium autumnale, 261; **scaposum**, 469
Helianthus annuus, 505, 506, 268; *divaricatus*, 268; *scaberrimus*, 269; *strumosus*, 268; *tuberosus*, 269
Heleochloa schoenoides, 626
Hemerocallis, 399
Hemieva ranunculifolia, 351
Hepatica triloba, 253
Heptanthus brevipes, 454, 455; *cochlearifolius*, 454; **cordifolius**, 454, 455; **lobatus**, 454, 455; *ranunculoides*, 454, 455; **Shaferi**, 454, 455
Heracleum candicans, 639; *pinnatum*, 639; *Thomsoni*, 639
Herminium Monorchis, 629
Heuchera flabellifolia, 359; *flavescens*, 361; *glabra*, 353; *grossularifolia*, 359; *ovalifolia*, 351; *parviflora*, 356
Hieracium austro-americanum, 320; *triste*, 320
 Hilgard, Botanical work of Dr. E. W., 389
Hippeastrum solandriflorum, 403
Hippophae rhamnoides, 638; *salicifolia*, 638
Hippurus vulgaris, 346, 638, 587
Homalobus Bourgovii, 358; *humilis*, 361
 HOOKER, H. D., JR. Physiological observations on *Drosera rotundifolia*, 1
Hordeum jubatum, 345; *nodosum*, 628; *vulgare*, 628
 HOWE, M. A. A note on the structural

- dimorphism of sexual and tetrasporic plants of *Galaxaura obtusata*, 621
 HOWE, M. A. & COLLINS, F. S. Notes on species of *Halymenia*, 169
*Hulsea carnos*a, 359
Hyalopora Polypodii, Observations on the fern rust, 195
Hydrocotyle americana, 242
 Hydrophyllaceae, 318

Hygrohypnum dilatatum, 239, 241; eugyrium, 239, 241; ochraceum, 239; **peruvienne**, 332
Hylocomium brevirostre, 240; splendens, 244
Hymenopappus cinereus, 362
Hymenostylium curvirostre, 240, 241
Hyoscyamus pusillus, 642
Hypocoum, 586, 633
 Hypericaceae, 638
Hypericum perforatum, 583, 638
Hypnum Schreberi, 244, 333

Ibidium Beckii, 408
Impatiens, 581; biflora, 261; brachycentra, 583, 638; pallida, 242; Thomsoni, 583, 638
 Index to American botanical literature, 57, 113, 161, 201, 277, 335, 393, 435, 495, 539, 591, 651
Inula Helenium, 268; barbata, 583, 645; Falconeri, 645; *Helenium*, 268; obtusifolia, 645; rhizocephaloides, 583, 645; Royleana, 645
 Iridaceae, 311, 629
Iris ensata, 629; kamaoensis, 629; stylosa, 399
Isolepis conifera, 445; junciformis, 442, 447; leucostachya, 445; vestita, 446
Isopyrum grandiflorum, 585, 632
Ivesia Gordonii, 349; utahensis, 350

Jaeschkea latiseppala, 640
Jubula pennsylvanica, 239
 Juglandaceae, 630
Juglans cinerea, 252; nigra, 505, regia, 630
 Juncaceae, 628, 310
 Juncaginaceae, 626
Juncoides arcticum, 348; arcuatum, 348; glabratum, 351; hyperboreum, 347; intermedium, 346; parviflorum, 346; spicatum, 346; subapetalum, 360
Juncus effusus, 257; Drummondii, 349; Hallii, 360; himalensis, 629; lampocarpus, 629; leucomelas, 629; membranaceus, 582, 629; Mertensianus, 349; nevadensis, 353; Parryi, 349; Regelii, 353; sphacelatus, 628; triglumis, 629
Juniperus, 581; communis, 587, 626; macrospora, 582, 626; Wallichiana, 626
Jurinea ceratocarpa, 583, 646; ceratocarpa depressa, 583, 646

Kallymenia reniformis, 181
Kalmia latifolia, 243; microphylla, 348
 Kentrophyta tegetaria, 350
Kirchneriella contorta, 473; **elongata**, 473
 KNUDSON, L. Cambial activity in certain horticultural plants, 533
Kobresia capillifolia, 628; macrantha, 628; pygmaea, 628; Royleana, 628; Royleana Kokanica, 628; schoenoides, 585, 628
Kochia odontoptera, 631; prostrata, 580, 631
Koeleria argentea, 627; cristata, 587, 627
Koelpinia linearis, 583, 646
 KUNKEL, L. O. Further studies of the orange rusts of *Rubus* in the United States, 559

Labatia aristata, 460
 Labiatae, 641
Lactuca, 589; decipiens, 584, 647; longifolia, 647; orientalis, 647; sativa, 647; Scariola, 584, 589, 647; tartarica, 647; undulata, 647
 Ladak, The flora of, 571, 625
Lagostis glauca, 583, 643; globosa, 643
Lamium album, 583, 642; amolexicaule, 588, 642; rhomboideum, 642
Lancea tibetica, 580, 581, 642
Laportea canadensis, 242, 259
Larix Lyallii, 350, 351
Larnalles, 105; dichotoma, 105; glauca, 105
Lathyrus altaicus, 637; arizonicus, 361; magellanicus, 316; sativus, 637
 Lauraceae, 297
Laurophyllum elegans, 285, 297; nervillosum, 298
Lavatera kashmiriana, 581, 583, 638
Ledum glandulosum, 348; groenlandicum, 347
Lecanora bermudeusis, 157; cinereocarnea, 157; cinereocarnea **athallina**, 157; cinereocarnea **rugosa**, 157; claronella, 157; glaucomodes conflectens, 157; pallida cancriformis, 157; subfusca, 158; varia, 158; varia symmicta, 158
Lecidea euporiza, 152; fuscorubescens, 152; revertens, 152; semiusta, 152; sphaeroides vacillans, 153
Leersia virginica, 257, 261
 Leguminosae, 315, 636
Lejeunea cavifolia, 240
Lemna gibba, 349; minor, 346, 587, 628
 Lemnaceae, 628
 Lentibulariaceae, 643
Leontopodium alpinum, 581, 583, 586, 644
Lepargyrea canadensis, 345
Lepidium apetalum, 633; capitatum, 586, 633; latifolium, 580, 633; ruderales, 588
Leptasea austromantana, 355; chrysan-

- tha, 361; *Hirculus*, 346; *tricuspidata*, 348
Leptobryum *Wilsonei*, 328
Leptodactylon *Nuttallii*, 349; *pungens*, 357
Leptodontium integrifolium, 324
Leptogium *diaphanum*; *marginellum*, 155; *tenuissimum*, 156; *tremelloides*, 156
Leptorhabdos *Benthiana*, 643
Leptotrema trypaneoides, 151
Leskea *abbreviata*, 74, 75; *gracillima*, 332
Lesquerella montana, 360; *parvula*, 360; *Wardii*, 363
Leuceria lanigera, 320; *multifida*, 320; *patagonica*, 320; *purpurea*, 320; *run-cianata*, 320
Leucodon *Draceanae*, 71
Leucoloma *Holstii*, 64
Leucolejeunea *clypeata*, 240
Leucojum *vernum*, 401
Lichens of *Bermuda*, *The*, 145
Ligusticum *filicinum*, 358, 359; *Porteri*, 61; *Thomsoni*, 581, 639
Lilaea *subulata*, 548, 554
Liliaceae, 310, 629
Lilium, 399; *auratum*, 506; *Martagon*, 505, 506; *philadelphicum*, 548, 549, 550
Limnorchis, 554
Limnorchis purpurascens, 360
Limnosella aquatica, 587, 642
Linaceae, 637
Lindbergia *abbreviata*, 74; *Austini*, 74; **haplocladioides**, 75; **pseudoleskeoides**, 75
Lindelofia *Benthami*, 581, 641
Linnaea *americana*, 345; *longiflora*, 352
Linum *perenne*, 637
Liriodendron *tulipifera*, 262
Lithophragma *bulbifera*, 355
Lithospermum *arvense*, 588, 641
Lloydia *serotina*, 346, 629
Loasa *argentina*, 317; *volubilis*, 317
Loasaceae, 317
Lolium *perenne*, 588, 628
Lonicera *canadensis*, 244; *asperifolia*, 583; 644; *caerulea*, 644; *glaucia*, 581, 583, 644; *heterophylla*, 644; *microphylla*, 583, 644; *Myrtillus*, 644; *rupicola*, 644; *spinosa*, 581, 586, 644
Lophocolea *bidentata*, 239; *minor*, 241
Lophozia *alpestris*, 244; *attenuata*, 244; *badensis*, 241
Lotus *corniculatus*, 581, 583, 588, 637
Lupinus *caespitosus*, 357; *monticola*, 358; *parviflorus*, 357; *pulcherrimus*, 358
Lutkea *pectinata*, 351
Luzula *chilensis*, 310
Luzuriago *marginata*, 310
Lychnis *apetala*, 582, 632; *brachypetala*, 585; *himalayensis*, 582, 632; *macro-rhiza*, 585, 632; *nutans*, 582, 632
Lycium *ruthenicum*, 642
Lycopodium *magellanicum*, 309; *obscurum*, 246; *Selago*, 244
Lycopis *arvensis*, 588, 641
Lyngbya *Birgei*, 482; *Hieronimusii*, 482
Lysiella *obtusata*, 346
Machaonia calcicola, 452, 453; *littoralis*, 452; *microphylla*, 452, 453; *pauciflora*, 452; *trifurcata*, 452, 453
MACKENZIE, K. K. Notes on *Carex*—X, 423; —XI, 601
Macronema *discoidea*, 350; *grindeli-folium*, 355
Macrotomia *Benthami*, 641; *perennis*, 641
Madia *sativa*, 321
Magnoliaceae, 314
Malapoenna *horrellensis*, 285, 299
Malcolmia *africana*, 634
Malpighia *Shaferei*, 456
Malva *verticillata*, 588, 638
Malvaceae, 638
Manihotites *georgiana*, 285, 295
Marrubium *lanatum*, 586, 641
Marsupella *emarginata*, 240, 241
Matricaria *chamomilla*, 321
Matthiola *odoratissima*, 634
Maviscus *cyperoides*, 441; *gracilis*, 441; *Sieberianus*, 441
Maytenus *magellanicus*, 316
MCALLISTER, F. The morphology of *Thallocarpus* *Curtisii*, 117
Meconopsis *aculeata*, 584, 633
Medeola *virginiana*, 253
Medicago *falcata*, 588, 636; *lupulina*, 588, 636; *sativa*, 588, 636
Melandrium *alpestris*, 314; *magellanicum*, 313
Melanotheca *aggregata*, 148; *cruenta*, 148
Melica *ciliata*, 627; *cupani*, 627
Melilotus *alba*, 588, 636; *officinalis*, 588, 636
Mentha *arvensis*, 642; *longifolium*, 588; *sylvestris*, 642
Menyanthes *trifoliata*, 346
Mertensia *alpina*, 361; *Bakeri*, 361; *Drummondii*, 362; *echioides*, 641; *lateriflora*, 361; *nivalis*, 361; *Parryi*, 361; *polyphylla*, 361; *stenoloba*, 359; *Tweedyi*, 359; *viridis*, 363; *viridula*, 361
Meteoriopsis *remotifolia*, 331
Metrosideros, 387; *polymorpha*, 387
Metzgeria *conjugata*, 240
Micractinium *pusillum*, 477, 479; **quadri-setum**, 479
Microphiale *lutea*, 152
Microthamnium *saproadelphum*, 64, 78; *tenellum*, 78
Micranthes *arguta*, 349; *aestivalis*, 354; *arnoglossa*, 357; *brachypus*, 361; *Ly-allii*, 353; *occidentalis*, 358; *rhomboid-*

- dea, 357; saximontana, 358; Rydbergii, 358
- Microula Benthami**, 641
- Mielichhoferia andina**, 328; bogotensis, 328; procerrima, 68; procumbens, 68; promontorii, 68
- Mikania Stevensiana**, 458
- Milium effusum**, 587, 626
- Mimulus Langsfordii**, 349; strictus, 642
- Mitracarpum depauperatum**, 467
- Mittenothamnium andicola**, 333; oxy-stegum, 333
- Mnium cuspidatum**, 246; hornum, 239, 246; ligulatum, 329; orthorrhynchum, 241; punctatum, 239
- Moehringia laterifolia**, 346; macrophylla, 346
- Moneses reticulata**, 354; uniflora, 346
- Monotropia Hypopitys**, 505
- Moraceae**, 291, 630
- Morphology of Thallocarpus Curtisii**, The, 117
- Morphology of the flowers of Zea Mays**, 127
- Morus alba**, 588, 630
- Moschopsis spathulata**, 319
- Muhlenbergia comata**, 349; filiculmis, 360; gracilis, 360; mexicana, 261; racemosa, 345; subalpina, 360; Wolfi, 360
- Muscaria adscendens**, 357; delicatula, 357; micropetala, 363
- Myrcia havanensis**, 285, 300
- Myrica ripleyensis**, 285, 288
- Myricaceae**, 288
- Myrtaceae**, 300
- Myricaria elegans**, 638; germanica, 638; germanica prostrata, 638
- Myurella julacea**, 241
- Myzodendraceae**, 313
- Myzodendron linearifolium**, 313; quadriflorum, 313; punctulatum, 313
- Naiadaceae**, 626
- Naiocrene parvifolia**, 353
- Nantucket**, The ferns and flowering plants of, —XVII, 265
- Nassauvia Darwinii**, 321; pygmaea, 321; revoluta, 321; subspinosa, 321
- Neckera eucarpa**, 331; Jamesoni, 331; Lindigii, 331; pennata, 240
- Nemexia biltmoreana**, 413; cerulea, 414; diversifolia, 417; ecirrhata, 419; herbacea, 414, 417; herbacea melica, 417; Hugeri, 420; lasioneuron, 417; nigra, 414; pulverulenta, 420; tamnifolia, 411; tenuis, 419
- Nepeta botryoides**, 641; discolor, 583, 641; eriostachya, 642; floccosa, 581, 642; glutinosa, 584, 642; leucolaena, 583, 642; linearis, 641; longibracteata, 585, 642; nivalis, 586, 642; salviaefolia, 642; Thomsoni, 641; tibetica, 586, 642
- Nephralles**, 100; parviflora, 100
- Nephrocytium allantoideum**, 475
- Nephrolepis biserrata furcans**, 215, 216, 230; cordifolia, 215, 217; davallioides, 216; Duffii, 215; exaltata, 207, 215–217, 224, 230; exaltata bostoniensis, 207–233; tuberosa, 217
- Nephrolepis exaltata bostoniensis**, sports of, 232
- Nephroma antarcticum**, 308; arcticum, 308
- New and rare African mosses**, from Mitten's herbarium and other sources, 63
- New or interesting algae from the lakes of Wisconsin**, 471
- New species of Opuntia**, 83
- New species of Pritchardia**, A, 385
- NICHOLS**, G. E. The vegetation of Connecticut—V. Plant societies along rivers and streams, 235
- Nicotiana rustica**, 642; Tabacum, 642
- Note on the structural dimorphism of sexual and tetrasporic plants of Galaxaura obtusata**, A, 621
- Notes on Carex**, —X, 423; —XI, 601
- Notes on plants of the southern United States**—I, 93; —II, 407
- Notes on species of Halymenia**, 169
- Nothofagus antarctica**, 306, 313
- Nymphaea polysepala**, 349
- Oakesia sessilifolia**, 252
- Observations on the fern rust Hyolopsora Polypodii**, 195
- Oedipodium australe**, 67; Griffithianum, 67
- Oenothera**, 207; Hilgardi, 390; odorata, 317
- Oleaceae**, 640
- Omphalaria cubana**, 154; lingulata, 154
- Omphalodes Thomsoni**, 641
- Onagraceae**, 317, 638
- Oncostylis fimbriata**, 446; vestita, 446
- Onoclea**, 366; sensibilis, 366, 368, 382, 261; Struthiopteris, 260
- Oocystis, panduriformis**, 471; panduriformis **minor**, 471
- Onosma echioides**, 641
- Opegrapha atra**, 149; Bonplandi, 149; chevalieri incarnata, 150; **ophites**, 150; vulgata, 150
- Ophioglossaceae**, 309
- Ophrys borealis**, 356; nephrophylla, 355
- Opuntia**, Additional species of, 523; New species of, 83
- Opuntia alta**, 84; **anahuacensis**, 92; basilaris, 83, 84; Bergeriana, 531; brachyclada, 84; **calantha**, 524; camanchica, 530; chlorotica, 89; cirbe, 524; **columbiana**, 523; **crystalenia**, 528; **cucumiformis**, 524; **curvospina**, 88;

- deltica*, 84; *flexospina*, 87; *Hanburyana*, 529; *ithypetala*, 529; *humistrata*, 83; *laxiflora*, 85; *littoralis*, 90; *longiclada*, 525; *megalantha*, 530; *microcarpa*, 527; *microdisca*, 525; *occidentalis*, 90; *platynoda*, 526; *pyrocarpa*, 90; *rubiflora*, 529; *semispinosa*, 89; *squarrosa*, 91; *zuniensis*, 86
- Orchidaceae, 312, 629
- Orchis latifolia*, 629; *rotundifolia*, 348
- Oreobroma minima*, 363
- Oreochrysum Parryi*, 362
- Oreoxis alpina*, 361; *Bakeri*, 361; *humilis*, 361
- Origanum vulgare*, 583, 642
- Origin of new species of *Nephrolepis* by orthogenetic saltation, The,—I. Progressive variations, 207
- Orobanchaceae, 643
- Orobanche cernua*, 643; *Hansii*, 643
- Orontium aquaticum*, 255
- Oryzopsis lateralis*, 626; *Munroi*, 626
- Osmorrhiza Berterii*, 317; *Leibergii*, 351, 352; *obtusata*, 357
- Osmunda Claytoniana*, 252
- Ostrya virginiana*, 252
- Ourisia rupestris*, 318
- Oxalidaceae, 316
- Oxalis acetosella*, 244; *patagonica*, 316
- Oxycoccus Oxycoccus*, 347
- Oxygraphis polypetala*, 632
- Oxypolis Fendleri*, 361
- Oxyria digyna*, 584, 587, 630
- Oxyrrhynchium rusciforme*, 239
- Oxytropis cachemirica*, 581, 637; *densa*, 586, 637; *lapponica*, 586, 637; *microphylla*, 586, 637; *mollis*, 637; *tatarica*, 586, 637; *Thomsoni*, 637
- Ozomelis Parryi*, 361; *stauropetala*, 355, 361; *trifida*, 351
- Pachyanthus reticulatus*, 459; *myrsinites*, 348
- Palicourea elongata*, 468
- Palamocladium leskeoides*, 333
- Palmae*, 288
- Panicularia nervata*, 345; *pauciflora*, 349
- Panicum agrostoides*, 257; *miliaceum*, 582, 588, 626
- Papaver nudicaule*, 633
- Papaveraceae, 633
- Papillaria imponderosa*, 331
- Paracaryum heliocarpum*, 641; *himalayense*, 641; *tibeticum*, 641; *Thomsoni*, 641
- Parietaria debilis*, 630
- Paris quadrifolia*, 400
- Parmelia latissima cristifera*, 158; *perlata*, 158; *tinctorum*, 158
- Parnassia fimbriata*, 349; *Kotzebuei*, 347; *ovata*, 635; *palustris*, 347, 635
- Paronychia pulvinata*, 360
- Parrya exscapa*, 634; *lanuginosa*, 634; *macrocarpa*, 634
- Paspalum ambiguum*, 626; *Curtisianum*, 95
- Paulownia tomentosa*, 43–56
- Pectianthia Breweri*, 353; *pentandra*, 349
- Pediastrum angulosum araneosum*, 476; *araneosum*, 476; *araneosum rugulosum*, 476; *Boryanum*, 476; *sculptatum*, 475
- Pedicularis bicornuta*, 583, 643; *bracteata*, 349; *cheilanthifolia*, 643; *dolichorrhiza*, 643; *Grayi*, 362; *Hallii*, 363; *longiflora*, 580, 581, 643; *lunata*, 363; *mollis*, 643; *Oederi*, 643; *Parryi*, 362; *pycnantha*, 643; *racemosa*, 349; *rhinanthoides*, 643; *scopulorum*, 363; *siphonantha*, 643; *tenuirostris*, 643
- Peganum Harmala*, 638
- Pellia epiphylla*, 239, 246; *Fabroniana*, 241
- Peltandra virginica*, 255
- PENNEL, F. W. Notes on plants of the southern United States, —I, 93; —II, 407
- Pennisetum flaccidum*, 580, 626; *lanatum*, 626
- Pentstemon aridus*, 359; *caespitosus*, 361; *crassifolium*, 351; *fruticosus*, 351; *Hallii*, 361; *Harbourii*, 361; *Lyallii*, 359; *montanus*, 359; *procerus*, 349; *pseudohumilis*, 359; *stenosepalus*, 361; *Tweedyi*, 359
- Perezia linearis*, 321; *recurvata*, 321
- Pernettya mucronata*, 317
- Peroniella Hyalothecae*, 477; *planctonica*, 476
- Perowskia abrotanoides*, 642
- Pertusaria communis lutescens*, 156; *leioplaca*, 156; *lutescens*, 156; *multi-punctata*, 157; *papillata*, 156; *pustulata*, 157; *tuberculifera*, 157
- Peruvian mosses, 323
- Petasites corymbosa*, 354; *frigida*, 354
- Phacelia alpina*, 359; *ciliolata*, 357; *circinata*, 318; *heterophylla*, 349; *Lyallii*, 359; *nervosa*, 361; *sericea*, 357
- Phaeographis lobata*, 151
- Phalaris arundinacea*, 262
- Phegopteris polypodioides*, 242, 244, 376, 382
- Philonotis fontana*, 239
- Phleum alpinum*, 582, 626; *pratense*, 345
- Phlomis bracteosa*, 583, 642
- Phlox alyssoides*, 357; *caespitosa*, 355; *condensata*, 361; *costata*, 358; *depressa*, 357; *diapensioides*, 358; *Kelseyi*, 357
- Phragmidium*, 566
- Phragmites communis*, 587, 627; *Phragmites*, 345; *Prattii*, 285, 287
- Phyllanthus nanus*, 464
- Phyllodoce empetriformis*, 348; *glanduliflora*, 351

- Physalis Alkekengi*, 642
Physcia alba, 160; *crispa*, 160; *integrata soresiosa*, 160; *Leana*, 160; *picta*, 160; *stellaris*, 160
 Physiological observations on *Drosera rotundifolia*, 1
Physochlaina praealta, 642
 Phytogeographical notes on the Rocky Mountain region—VI. Distribution of the subalpine plants, 343
Picea albertiana, 355; *canadensis*, 354; *Engelmannii*, 354, 355; *mariana*, 354
Pilea pelpoides, 630; *pumila*, 242, 261
Pilosella Richardsoniana, 353
Pilotrichella conferta, 71; *viridis*, 331
Pimpinella Saxafraga, 588, 639
 Pinaceae, 626
Pinguicula vulgaris, 347
Pinus, 198; *albicaulis*, 348, 352; *aristata*, 350; *Balfouriana*, 350; *flexilis*, 348; *Murrayana*, 348; *palustris*, 93; *Strobus*, 506
Pisum sativum, 637
Pituranthos Thomsoni, 639
Plagiochila asplenioides, 239, 240
Plagiothecium denticulatum, 240; **nitens**, 78; *Mildbraedii*, 78
Plagiopus Oederi, 240, 241
 Plantaginaceae, 319, 643
Plantago brachyphylla, 643; *major*, 587, 643; *maritima*, 319; *tibetica*, 643
 Plants from southern Patagonia collected by Charles Wellington Furlong, 305
Platanus occidentalis, 252, 259, 262; *orientalis*, 39, 45, 56
Pleea tenuifolia, 408
Pleurogyne brachyanthera, 640; *carinthiaca*, 583, 640; *fontana*, 357; *Thomsoni*, 581; 640
Pleurospermum Candollii, 583, 639; *Hookeri*, 639; *stellatum*, 586, 639
Pluchea camphorata, 265
 Plumbaginaceae, 317, 640
Poa alpica, 360; *alpina*, 587, 627; *annua*, 582, 588, 627; *aperta*, 360; *argentina*, 310; *attenuata*, 627; *Buckleyana*, 356; *bulbosa*, 627; *crocata*, 345; *confusa*, 358; *Cusickii*, 352; *epilis*, 356; *Fenderleriana*, 349; *fuegianus*, 309; *Grayana*, 360; *laevigata*, 356; *leptocoma*, 355; *Lettermanii*, 360; *longiligula*, 349; *longipedunculata*, 360; *lucida*, 360; *nemoralis*, 582, 587; *nemoralis ligulata*, 585, 627; *nervosa*, 358; *nevadensis*, 355; *occidentalis*, 360; *Olneyi*, 356; *Paddensis*, 355; *Pattersonii*, 360; *pratensis*, 585, 587, 627; *Pritchardi*, 310; *pudica*, 363; *reflexa*, 355; *rupicola*, 355; *Sheldoni*, 360; *tibetica*, 627; *Traceyi*, 360; *tremula*, 627; *triclepis*, 363; *Vaseyana*, 360
 Poaceae, 287
Podophyllum Emodi, 581, 582, 633
Pogonatum campylocarpum, 330; *polycarpum*, 330; *simense*, 69
 Polygonaceae, 630
Polygonatum biflorum, 253; *verticillatum*, 629
 Polemoniaceae, 318
Polemonium Brandegei, 361; *confertum*, 361; *delicatum*, 361; *foliosissimum*, 361; *Grayanum*, 363; *miletum*, 361; *viscosum*, 359
Polygonum affine, 582, 584, 630; *alatum*, 631; *amplexicaule*, 630; *aviculare*, 580, 587, 630; *cognatum*, 584, 630; *Convolutus*, 631; *filicaule*, 630; *Hydropiper*, 587, 631; *interruptus*, 310; *islandica*, 630; *lapathifolium*, 630; *molliaeforme*, 584, 630; *Muhlenbergii*, 255; *paronychioides*, 630; *pennsylvanicum*, 257; *Persicaria*, 631; *polystachyum*, 582, 631; *scandens*, 257, 260; *sibiricum*, 585, 631; *sphaerostachyum*, 630; *tortuosum*, 582, 585, 631; *tubulosum*, 630; *vaccinifolium*, 630; *viviparum*, 587, 630
 Polypodiaceae, 309, 625
 Polypodiaceae, Branched prothallia in the, 365
Polypodium, 210; *aureum*, 365, 377; *vulgare*, 242
Polypogon monspeliensis, 588, 627
 Polystichum, 210
Polytrichadelphus magellanicus, 309
Polytrichum alpinum, 244; *antillarum*, 330
Pontederia cordata, 255
Populus, 589; *alba*, 588, 629; *ciliata*, 629; *candicans*, 588, 629; *deltoides*, 259; *euphratica*, 629; *nigra italica*, 588, 629; *tremuloides*, 344
Porella pinnata, 239; *platyphylloides*, 240
Porina nucula, 147; *phaea*, 147; *tetracerae*, 147
Powtrichum valdiviae, 331
Porphyra amplissima, 171
Potamogeton alpinum, 345; *filiformis*, 345; *heterophyllum*, 345; *interior*, 356; *nalus*, 345; *Richardsonii*, 345; *pectinatus*, 345, 585, 587, 626; *perfoliatus*, 587, 626; *sp.*, 255
Potentilla ambigua, 586, 635; *anserina*, 315, 580, 587, 635; *arachnoides*, 357; *argyrophylla*, 582, 584, 636; *atro-sanguinea*, 636; *bifurca*, 580, 581–586; *curviseta*, 635; *divisa*, 357; *flabellifolia*, 353; *fruticosa*, 635; *fruticosa Inglisii*, 582, 584, 587, 635; *fruticosa ochreata*, 635; *fruticosa pumila*, 635; *filipes*, 357; *gelida*, 636; *glaucochylla*, 357; *glomerata*, 358; *Macounii*, 358, 359; *minutifolia*, 363; *modesta*, 361; *multifida*, 581, 586, 635; *Nelsonii*, 363; *nivea*, 346, 636;

- ovina, 358; paucijuga, 363; perdissecta, 358; quinquefolia, 346; saximontana, 357; sericea, 587, 635; Salesovii, 635; Sibbaldi, 582, 635; tenerrima, 363; tetrandra, 587, 635; uniflora, 346; virgulata, 363; viridis, 361; wyomingensis, 363
- Prangos pabularia, 639
- Preissia quadrata, 241
- Primula angustifolia, 361; brevicalyx, 639; denticulata capitata, 639; elliptica, 639; farinosa, 587, 639; Heydei, 639; involucrata, 639; Meeboldii, 639; Parryi, 357; purpurea, 639; sibirica, 581, 586, 639; Stuartii, 583, 639
- Primulaceae, 317, 639
- Prionodon bolivianus, 331
- Pritchardia arecina, 385; **Beccariana**, 385, 386; eriophora, 385; eriostachya, 385; Gaudichaudii, 385; Hillebrandii, 385; lanigera, 385; Martii, 385; minor, 385; remota, 385; Rockiana, 385
- Pritchardia, A new species of, 385
- Proteaceae, 313
- Prunus armeniaca, 636; persica, 636; tomentosa, 636
- Psedra quinquefolia, 260
- Pseudocymopterus montanus, 361; purpureus, 354; sylvaticus, 361; Tidestromii, 361
- Pseudoleskea denudata, 77; dispersa, 76
- Pseudoreoxys bipinnata, 358
- Pseudopteryx anisata, 361; longiloba, 361
- Pseudotsuga mucronata, 348, 355
- Psorotichia **bermudana**, 154; diffracta, 154
- Pteridium, 366; aquilinum, 366, 380, 382
- Pterigynandrum filiforme, 241
- Pterospermites carolinensis, 285, 297
- Ptilium Crista-castrensis, 244
- Puccinia Peckiana, 559, 563
- Pulsatilla occidentalis, 353
- Pyrenula aurantiaca, 147; brachysperma, 147; leucoplaca, 148; manillana, 148; nitida nitidella, 148
- Pyrgillus cubanus, 148
- Pyrola chlorantha, 346; minor, 346; secunda, 346; uliginosa, 346; rotundifolia, 246
- Pyrrocoma Clementis, 362
- Pyrus communis, 635; Malus, 589, 635
- Quantitative study of the factors influencing the weight of the bean seed—II. Correlation between the number of pods per plant and seed weight, 485
- Quercus bicolor, 262; palustris, 262; rubra, 252
- Quinchamalium procumbens, 313
- Racomitrium aciculare, 239, 241
- Radicula alpina, 356; curvipes, 363; palustris, 257; Underwoodii, 363
- Radula complanata, 240; obconica, 241
- Rajania cordifolia, 546; hastata, 546
- Ramalina complanata, 158
- Ranunculaceae, 293, 314, 632
- Ranunculus adoneus, 360; affinis, 356, 633; alismaefolius, 360; alpeophilus, 356; aquatilis, 587, 633; Cymbalaria, 314, 580, 587, 632; eximius, 356; hirtellus, 582, 584, 632; Helleri, 358; hyperboreus, 347, 633; laetus, 633; Macauleyi, 360; micropetalus, 360; peduncularis, 314; pulchellus, 580, 582, 585, 632; saxicola, 358; stenolobus, 360; Suksdorfii, 351
- Raphidium setigerum, 473
- Reinschiella setigera, 473
- Relbunium pusillum, 319
- Rhabdoweisia denticulata, 240, 241
- Rhacomitrium crispipilum, 328
- Rhacopilum tomentosum, 332
- Rhamnaceae, 316
- Rheum Emodi, 630; spiciforme, 585, 630; tibeticum, 630; Webbianum, 630
- Rhodiola polygama, 361
- Rhus Toxicodendron, 260
- Rhynchostegium aquaticum, 334; conophyllum, 334
- Rhynchostegiella Holstii, 80; Volkensii, 80
- Ribes coloradensis, 359; cucullatum, 315; glandulosum, 352; Grossularia, 588, 635; hudsonianum, 347; laxiflorum, 352; montigenum, 348; orientale, 582, 635; parvulum, 356; petiolaris, 351; Wolfii, 359
- Richteriella botryoides, 478, 479; globosa, 478; quadriseta, 479
- Riccia crystallina, 120; Curtisii, 117, 119; Frostii, 121; glauca, 122
- RIDDLE, L. W. The lichens of Bermuda, 145
- Rigodium **dentatum**, 64, 79; gracile, 79; kilimandscharico, 79
- Rinodium insperata, 159
- Rochelia cardiosepala, 641; rectipes, 641; stellulata, 641
- ROCK, J. F. A new species of Pritchardia, 385
- Romanzoffia Leibergii, 363; sitchensis, 353
- Rondeletia **calicicola**, 467
- Rosa Eglanteria, 636; Webbiana, 581, 582, 636
- Rosaceae, 315, 635
- ROWLEE, W. W. Plants from southern Patagonia collected by Charles Wellington Furlong, 305
- Rubia tibetica, 581, 586
- Rubiaceae, 643
- Rubus acaulis, 347; americanus, 567; arcticus, 567; argutus, 563; canadensis,

- 561, 564, 567; *Chamaemorus*, 347; *cuneifolius*, 567; *frondosus*, 561, 567; *nigrobaccus*, 561, 567; *nivalis*, 351; *occidentalis*, 567; *procumbens*, 564; *Randii*, 567; *saxatilis*, 559, 567, 635; *stellatus*, 567; *strigosus*, 567; *triflorus*, 567; *trivialis*, 567; *vermontanus*, 567; *villosus*, 567; *vitifolius*, 567
- Rudbeckia hirta*, 268
- Rumex Acetosa*, 588, 630; *orientalis*, 630
- RYDBERG, P. A. Phytogeographical notes on the Rocky Mountain region—VI. Distribution of the subalpine plants, 343
- Rydbergia Brandegei*, 362; *grandiflora*, 357
- Rynchospora intermixta*, 443; *oligantha*, 443; *pusilla*, 443
- Sabalites* sp., 285, 288
- Saelania glaucescens*, 241
- SAFFORD, W. E. *Desmopsis*, a new genus of *Annonaceae*, 183
- Sagina procumbens*, 587; *saginoides*, 346
- Sagittaria angustifolia*, 95; *latifolia*, 255; *variabilis*, 548, 554
- Salicaceae*, 289, 629
- Salix alaxensis*, 353; *alba*, 588, 629; *angustifolia*, 629; *Barclayi*, 351; *casadensis*, 352; *chlorophylla*, 345; *cordata*, 258; *daphnoides*, 582, 629; *desertorum*, 353; *divergens*, 630; *Drummondiana*, 353; *elegans*, 629; *eutawensis*, 285, 289; *Fernaldii*, 358; *flabellaris*, 629; *fragilis*, 588, 629; *glaucops*, 348; *hastata*, 629; *longifolia*, 258; *nigra*, 259; *oxycarpa*, 630; *myrtillifolia*, 353; *pseudomyrsinites*, 347; *pycnostachya*, 629; *saskatchewanana*, 353; *saximontana*, 356; *sclerophylla*, 629; *Seemanii*, 353; *subcaerulea*, 351; *tetrasperma*, 630; *Tweedyi*, 358; *Wolfii*, 359
- Sambucus canadensis*, 260; *microbotrys*, 359
- Salsola collina*, 585, 631; *Kali*, 587, 631
- Samolus spathulatus*, 317
- Sanguinaria canadensis*, 253
- Sanicula gregaria*, 242
- Santalaceae*, 313, 630
- Saponaria Vaccaria*, 632
- Satureia Darwinii*, 319
- Saussurea albescens*, 646; *bracteata*, 646; *elliptica*, 646; *Falconeri*, 646; *glanduligera*, 646; *graminifolia*, 646; *Jacea*, 646; *Schultzii*, 646; *sorocephala*, 646; *Stoliczkai*, 646; *subulata*, 646; *taraxifolia*, 646; *Thomsoni*, 646
- Saussurea*, 589; *albescens*, 583; *bracteata*, 586; *Jacea*, 583; *glanduligera*, 586; *sorocephala*, 583
- Savastana odorata*, 345
- Savia perlucens*, 464
- SAX, K. Fertilization in *Fritillaria pudica*, 505
- Saxifraga*, 361, 589; *cernua*, 346, 582, 635; *cordillerarum*, 315; *debilis*, 356; *flagellaris*, 582, 635; *Hirculus indica*, 582, 635; *Jacquemontiana*, 635; *oppositifolia*, 587, 635; *ramulosa*, 635; *rivularis*, 347; *sibirica*, 580, 584, 635; *Stracheyi*, 582, 635
- Saxifragaceae*, 314, 635
- Scapania memorosa*, 240, 241
- Scheuzeria palustris*, 345
- Schizmus marginatus*, 627
- Schoenocaulon Drummondii*, 408, 409; *texanum*, 408, 409
- Schoenocaulon* A. Gray in southern Texas, 408
- Schoenus paradoxus*, 444
- Schoepfia cubensis*, 458
- Schroederia Judayi*, 474; *setigera*, 473
- Scirpus americanus*, 255; *caespitosus*, 347; *Caricis*, 628; *capillaris*, 447; *cyperoides*, 441; *coarctatus*, 447; *floccosus*, 445; *hirtus*, 446; *juncoides*, 442, 447; *lacustris*, 628; *nodulosus*, 442; *pauciflorus*, 587, 628; *pumilus*, 628; *rufus*, 587, 628; *setaceus*, 628; *stenophyllus*, 443; *validus*, 255
- Scleria gracilis*, 443
- Sclerophyton elegans*, 151
- Scolopendrium*, 366; *vulgare*, 365, 378, 382
- Scorzonera divaricata*, 581, 647; *purpurea*, 647
- Scrophularia dentata*, 642; *lucida*, 642; *scabiosaefolia*, 583, 642; *variegata*, 642
- Scrophulariaceae*, 318, 642
- Scutellaria Heydeli*, 641; *prostrata*, 583, 641
- Sebdenia heteronema*, 171, 172
- Sebesten brachycalyx*, 457; *Sebestena*, 457
- Sedum*, 589; *asiaticum*, 582, 635; *Ewersii*, 582, 635; *quadrifidum*, 635; *Rhodiola*, 582, 584, 587, 635; *tillaeoides*, 635; *tibeticum*, 582, 584, 586, 635
- Sematophyllum carolinianum*, 241
- Sempervivum acuminatum*, 580, 582, 586, 635
- Senecio admirabilis*, 362; *alpicola*, 359; *acanthifolius*, 321; *ambrosioides*, 362; *amplectens*, 362; *anacletus*, 362; *arnicoides frigida*, 583, 586, 646; *atratus*, 362; *carthamoides*, 362; *chloranthus*, 362; *chrysanthemoides*, 583, 646; *coronopifolius*, 646; *crassulus*, 362; *crocatus*, 362; *cymbalarioides*, 357; *Darwinii*, 321; *dubius*, 646; *falklandicus*, 321; *foliosus*, 362; *Harbourii*, 362; *Holmii*, 362; *Hatcherianus*, 321; *Jonesii*, 362; *Kingii*, 321; *lapathifolium*, 362; *longipes*, 321; *lugens*, 354; *megacephalus*, 359; *multicapitatus*, 362; *Nelsonii*, 362;

- paeninsularis, 321; pagosanus, 362;
 pauciflorus, 347; pedunculatus, 646;
 perplexans, 362; pentadontus, 362;
 pudica, 362; Purshianus, 357; pseu-
 dareus, 357; sericeo-nitens, 321; serido-
 phyllus, 350; soldanella, 362; sphaero-
 cephalus, 359; taraxacoides, 362; tibeti-
 cus, 646; trifurcatus, 321; vulgaris, 274;
 werneriaefolius, 362
Setaria viridis, 588, 626
Sibbaldia procumbens, 346
Sicyos angulatus, 260
Sieversia ciliata, 355
Silene, 358; *acaulis*, 346; *conoidea*, 632;
 Douglasii, 356; *Hallii*, 360; *Menziesii*,
 356; *Moorcroftiana*, 580, 584, 632;
 oregana, 351; *repens*, 353; *tenuis*, 582,
 632; *vulgaris*, 582, 632
Sisymbrium, 589; *Columnnae*, 580, 633;
 humile, 633; *mollissimum*, 633; *minuti-*
 florum, 633; *Sophia*, 633; *Thalianum*, 633
Sisyrinchium alpestre, 360; *chilense*, 311;
 iridifolium, 311, 399; *juncum*, 311;
 striatum, 311
Sitanion rigidum, 358
Smelkowskia americana, 356; *lineariloba*,
 360; *lobata*, 358; *ovalis*, 351
Smilacina racemosa, 242, 253
Smilax biltmoreana, 411, 413; *diversifolia*,
 417; *ecirrhata*, 411, 419; *herbacea*, 410,
 411, 413, 414; *Hugerii*, 411, 420; *inermis*,
 414; *lasioneuron*, 410, 411, 417;
 leptanthera, 410, 412; *peduncularis*,
 414; *Pseudo China*, 410, 413; *pulveru-*
 lenta, 410, 411, 420; *tamnifolia*, 410,
 411; *tenuis*, 411, 419
Smilax, Subgenus *Nemexia* (Raf.) in the
 eastern United States, 409
 SMITH, G. M. New or interesting algae
 from the lakes of Wisconsin, 471
 SMITH, P. M. The development of the
 embryo and seedling of *Dioscorea*
 villosa, 545
 Solanaceae, 642
Solanum Dulcamara, 260; *nigrum*, 642;
 tuberosum, 642
Solenanthus circinatus, 641
Solenomelus Sisyrinchium, 311
Solidago ciliosa, 357; *decumbens*, 362;
 latifolia, 242; *oreophila*, 357; *scopu-*
 lorum, 358; *serotina*, 261
Sonchus asper, 647; *oleraceus*, 589, 647
Sophia purpurascens, 360; *sophicoides*, 353
Sophora alopecuroides, 636; *Moorcrof-*
 tiana, 636
Sparganium, 554; *angustifolium*, 345;
 minimum, 345
Spathularia Brunnoniana, 354; *ferru-*
 ginea, 354; *Vreelandii*, 357
Sphaerocarpos, 117, 119; *californicus*
 119; *cristatus*, 123; *Donnellii*, 123;
 texanus, 119
Sphaerostigma Hilgardi, 390
Sphagnum, 2
Spermacoe havanensis, 453; *spinosa*, 453
Sphenolobus exsectus, 241
Spirulina laxa, 481
Spraguea multiceps, 358
Squamidium rotundifolium, 331
Stachys tibetica, 581, 642
Statice macrorrhachis, 640
Staurogenia apiculata, 472
Stellaria debilis, 314; *decumbens*, 584,
 631; *graminea*, 582, 585, 588, 631;
 glauca, 588, 631; *media*, 588, 631;
 semivestita, 632; *tibetica*, 631
Stenophyllum alpestris, 44, 446; *antil-*
 lanus, 444, 447; *capillaris*, 444, 447;
 ciliatifolius, 95; *coarctatus*, 444, 447;
 coniferus, 444, 445; *curassavicus*, 445;
 fimbriatus, 444, 446; *floccosus*, 444,
 445; *junciformis*, 442, 444, 447; *leuco-*
 stachys, 444, 445; *paradoxus*, 444;
 portoricensis, 444, 445; *subaphyllus*,
 443, 444, 446; *tenuifolius*, 448; *Tuerck-*
 heimii, 444, 446; *vetistus*, 444, 446;
 Wilsoni, 444, 446
 Sterculiaceae, 296
Sterculia Snowii tennesseensis, 285, 296
Stereodon cupressiformis, 246
Stereophyllum odontocalyx, 73
 STEWART, R. R. The flora of Ladak,
 western Tibet—I. Discussion of the
 flora, 571; —II. Check list of Ladak
 plants, 625
Sticta endochrysea, 308
Stipa barbata, 585, 626; *basiplumosa*, 626;
 Hookeri, 626; *minor*, 356; *mongolica*,
 626; *oregonensis*, 352; *Nelsonii*, 356;
 pogonathera, 310; *purpurea*, 626; *pur-*
 purea longe-aristata, 626; *sibirica*, 626;
 splendens, 626; *Tweedyi*, 356
Stracheya tibetica, 637
Streptopus amplexifolius, 244; *roseus*, 242,
 244
 Studies of West Indian plants—VIII, 441
Suaeda corniculata, 631; *microsperma*,
 631
Swertia congesta, 357; *cordata*, 583, 640;
 petiolata, 583, 640; *scopulina*, 357
Symphystemon biflorus, 311; *Lyck-*
 holmi, 311
Synthyris dissecta, 359; *laciniata*, 359;
 pinnatifida, 363; *reniformis*, 351, 352
 Tamaricaceae, 638
Tamarix gallica Pallasii, 638
Tamus communis, 545, 546, 548, 550,
 551, 555
Tanacetum, 589; *artemisioides*, 583, 645;
 fruticosum, 586, 645; *gracile*, 645;
 longifolium, 645; *nanum*, 645; *tibeti-*
 cum, 645; *tomentosum*, 645; *vulgare*,
 273

- Tapura obovata*, 465
Taraxicum laevigatum, 32; *leucanthum*, 647; *officinale*, 581, 584, 586, 589, 647
Tauscheria lasiocarpa, 634
Telesonix heucheriformis, 358; *Jamesii*, 361
Tennessee, Contributions to the Mesozoic flora of the Atlantic coastal plain, 283
Tetraedron enorme, 480; *gracile*, 480; *limneticum*, 480; *lobatum polyfurcatum*, 480; *planctonicum*, 479
Tetrastrum apiculatum, 472
Thalictrum, 589; *alpinum*, 346, 633; *dioicum*, 253; *minus*, 633; *minus foetidum*, 633; *platycarpum*, 633; *rutaefolium*, 633; *venulosum*, 356
Thallocarpus Curtisii, The morphology of, 117
Thamnum alleghaniense, 240; *alopecurum*, 71; *capense*, 71; *Leibergii*, 71; *maderense*, 71
Thelidium bermudanum, 146; *Farlowi*, 147; *pyrenophorum*, 147
Thelotrema trypaneoides, 141
Thermometric movements of tree branches at freezing temperatures, The, 29
Thermopsis inflata, 582, 636
Thesium himalense, 630
Thlaspi arvense, 582, 633; *alpestre*, 633; *californicum*, 353; *coloradense*, 360; *glaucum*, 356; *magellanicum*, 314; *purpurascens*, 360
Thuidium delicatulum, 239; *peruvianum*, 332
Thylacospermum rupifragum, 632
Thymus Serpyllum, 586, 588, 642
Tilia americana, 252, 260, 262; *europaea*, 29
Tofieldia intermedia, 351
Tonestus laceratus, 363; *pygmaeus*, 362
Tortella tortuosa, 241
Tortula affinis, 326; *andicola*, 326; *confusa*, 326; *erythroneura*, 67; *Eubryum*, 66; *exesa*, 67; *lacerifolia*, 326; *laevipila*, 66; *pagorum*, 66; *ruralis*, 67
Tournefortia Earlei, 460; *incana*, 460; *stenophylla*, 460
Townsendia grandiflora, 362
Trachydium Roylei, 583, 639
Trachyphyllum fabronioides, 72; *gastrodes*, 72
Tragopogon porrifolius, 647; *pratense*, 583, 589, 646
Tribulus terrestris, 588, 638
Tricholepis tibetica, 646
Trichopus zeylanicus, 546, 548
Trichostomum afrum, 66; *afro-fontanum*, 66; *brachydontium*, 324
Trifolium attenuatum, 363; *bracteolatum*, 363; *Brandegei*, 361; *dasyphyllum*, 361; *Haydeni*, 358; *lividum*, 361; *montanense*, 358; *nanum*, 357; *Parryi*, 361; *pratense*, 588, 636; *scariosum*, 363; *stenolobum*, 361
Triglochin maritima, 345, 580, 585, 587, 626; *palustris*, 345, 580, 587, 626
Trigonella corniculata, 636; *Emodi*, 636
Trigyneia, 184; *Galeottiana*, 184, 187
Trillium erectum, 242, 253
Trisetum majus, 355; *montanum*, 360; *subspicatum*, 345, 582, 627
Triticum vulgare, 628
Trollius albiflorus, 355
TROWBRIDGE, C. C. The thermometric movements of tree branches at freezing temperatures, 29
Tsuga Mertensiana, 352
Tium alpinum, 346
Tulipa Gesneriana, 400, 548
Tussilago Farfara, 589, 646
Typha latifolia, 255, 261, 262
Ulmaceae, 630
Ulmus americana, 252; *parvifolia*, 630
Ulotia americana, 240, 241
Umbelliferae, 317, 639
Undescribed species from Cuba, 458
Undescribed species from Porto Rico, 456
Undescribed species from the Isle of Pines, 462
Unona, 183; *hibracteata*, 183, 184, 190; *Desmos*, 183; *discolor*, 183; *discreta*, 183; *panamensis*, 183-185; *violacea*, 190
Unonopsis, 183, 184; *angustifolia*, 184; *Perottetii*, 184; *rufescens*, 184
Uredinopsis, 198
Urtica dioica, 588, 630; *hyperborea*, 585, 630; *magellianica*, 313
Urticaceae, 313, 630
Utricularia minor, 346, 587, 643; *vulgaris*, 346
Vaccinium caespitosum, 345; *globularis*, 359; *occidentale*, 352; *oreophilum*, 356, 357; *scoparium*, 356, 357; *uliginosum*, 347; *Vitis-Idaea*, 347
Vagnera stellata, 346
Valeriana acutiloba, 362; *carnosa*, 319; *dioica*, 644; *edulis*, 357; *septentrionalis*, 347; *sitchensis*, 354; *lapathifolia*, 319; *Stracheyi*, 644; *Wallichii*, 644
Valerianaceae, 319, 644
Vallisneria spiralis, 255
Vegetation of Connecticut, The,—V. Plant societies along rivers and streams, 235
Veratrum reciosum, 349
Verbascum Thapsus, 583, 588, 642
Verbena Carroo, 319; *hastata*, 261; *Pritchardi*, 319
Veronica agrestis, 589, 643; *americana*, 346; *Anagallis-aquatica*, 587, 643; *Beccabeenga*, 588, 643; *biloba*, 643;

- ciliata*, 586, 642; *deltigera*, 583, 642; *scutellata*, 347; *serpyllifolia*, 346; *Wormskioldii*, 346
Verrucaria albido-atra, 146; *bermudana*, 145, 146; *runderella*, 146; *rupestris*, 146; *rupestris ruderum*, 146
Vesicularia sphaerocarpa, 79
Viburnum acerifolium, 243
Vicia Faba, 637; *magellanica*, 316; *sericella*, 316; *tenuifolia*, 637
Viola adunca, 351; *bellidifolia*, 361; *blanda*, 242; *cinerea*, 638; *cucullata*, 242, 261; *flavovirens*, 358; *kunawarensis*, 638; *maculata*, 316; *neomexicana*, 361; *pubescens*, 253; *rotundifolia*, 242, 244
 Violaceae, 316, 638
 Vitaceae, 296
Vitis barbrusca, 533
Wahlberdella attenuata, 363; *Kingii*, 363; *montana*, 360
 WEATHERWAX, P. Morphology of the flowers of *Zea Mays*, 127
Werneria nana, 587, 646; *nana affinis*, 584, 646
Weymouthia mollis, 309
 WILLIAMS, R. S. Peruvian mosses, 323
Woodsia, 366; *obtusa*, 378, 382
Woodwardia, 366; *virginica*, 379, 382
 WUIST, E. D. Branched prothallia in the Polypodiaceae, 365
Xerophyllum tenax, 352
Xylopia frutescens, 183; *salicifolia*, 183
Xyris indica, 401; ***longibracteata***, 462
Yucca glauca, 408; sp., 408
Zamia silicea, 462
Zannichellia, 554; *palustris*, 587, 626
Zea Mays, Morphology of the flowers of, 127
Zizania palustris, 255
 Zygomphyllaceae, 638